

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 31/42, 31/425, 31/44, 31/505	A1	(11) International Publication Number: WO 94/25026 (43) International Publication Date: 10 November 1994 (10.11.94)
--	----	--

(21) International Application Number: PCT/GB94/00879

(22) International Filing Date: 25 April 1994 (25.04.94)

(30) Priority Data:
9308487.9 23 April 1993 (23.04.93) GB

(71) Applicant (for all designated States except US): SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): CAWTHORNE, Michael, Anthony [GB/GB]; SmithKline Beecham Pharmaceuticals, Great Burgh, Yew Tree Bottom Road, Epsom, Surrey KT18 5XQ (GB). HINDLEY, Richard, Mark [GB/GB]; SmithKline Beecham Pharmaceuticals, Great Burgh, Yew Tree Bottom Road, Epsom, Surrey KT18 5XQ (GB).

(74) Agent: RUTTER, Keith; SmithKline Beecham plc, Corporate Intellectual Property, SB House, Great West Road, Brentford, Middlesex TW8 9BD (GB).

(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KG, KP, KR, KZ, LK, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).


Published

With international search report.

(54) Title: USE OF THIAZOLIDINEDIONES FOR THE TREATMENT OF ATHEROSCLEROSIS AND EATING DISORDERS

(57) Abstract

The use of a compound of formula (I) or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, wherein: A¹ represents a substituted or unsubstituted aromatic heterocyclyl group; R¹ represents a hydrogen atom, an alkyl group, an acyl group, an aralkyl group, wherein the aryl moiety may be substituted or unsubstituted, or a substituted or unsubstituted aryl group; R² and R³ each represent hydrogen, or R² and R³ together represent a bond; A² represents a benzene ring having in total up to five substituents; and n represents an integer in the range of from 2 to 6, for the manufacture of a medicament for use in the treatment and/or prophylaxis of atherosclerosis and/or for the regulation of appetite and food intake.



FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

USE OF THIAZOLIDINEDIONES FOR THE TREATMENT OF ATHEROSCLEROSIS AND EATING DISORDERS

This invention relates to novel use of certain substituted thiazolidinedione derivatives and of pharmaceutical compositions containing such compounds.

- 5 European Patent Applications, Publication Numbers 0008203, 0139421, 0155845, 0177353, 0193256, 0207581 and 0208420 relate to thiazolidinedione derivatives which are disclosed as having hypoglycaemic and hypolipidaemic activity. Chem. Pharm. Bull 30 (10) 3580-3600 also relates to certain thiazolidinedione derivatives having hypoglycaemic and hypolipidaemic activities.
- 10 European Patent Application, Publication Number 0306228 discloses certain substituted thiazolidinedione derivatives of formula (A):




(A)

- 15 or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof, and/or a pharmaceutically acceptable solvate thereof, wherein:
- A^{1a} represents a substituted or unsubstituted aromatic heterocyclyl group;
- R^{1a} represents a hydrogen atom, an alkyl group, an acyl group, an aralkyl group,
- 20 wherein the aryl moiety may be substituted or unsubstituted, or a substituted or unsubstituted aryl group;
- R^{2a} and R^{3a} each represent hydrogen, or R^{2a} and R^{3a} together represent a bond;
- A^{2a} represents a benzene ring having in total up to five substituents; and
- n represents an integer in the range of from 2 to 6. Such compounds are disclosed
- 25 *inter alia* as being useful for the treatment and/or prophylaxis of cardiovascular disease and certain eating disorders.

- It has now surprisingly been discovered that these compounds are of particular use in the treatment and/or prophylaxis of atherosclerosis. In addition these compounds are particularly useful for the regulation of appetite and food intake in
- 30 subjects suffering from disorders associated with under-eating, such as anorexia nervosa, and disorders associated with over-eating, such as obesity and anorexia bulimia.

Accordingly, the present invention provides the use of a compound of formula

(I):



(I)

5 or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof, and/or a pharmaceutically acceptable solvate thereof, wherein:

A¹ represents a substituted or unsubstituted aromatic heterocyclyl group;

R¹ represents a hydrogen atom, an alkyl group, an acyl group, an aralkyl group, wherein the aryl moiety may be substituted or unsubstituted, or a substituted or

10 unsubstituted aryl group;

R² and R³ each represent hydrogen, or R² and R³ together represent a bond;

A² represents a benzene ring having in total up to five substituents; and

n represents an integer in the range of from 2 to 6, for the manufacture of a medicament for treatment and/or prophylaxis of atherosclerosis and/or for the

15 regulation of appetite and food intake in subjects suffering from disorders associated with under-eating, such as anorexia nervosa, and disorders associated with over-eating, such as obesity and anorexia bulimia.

Suitable aromatic heterocyclyl groups include substituted or unsubstituted, single or fused ring aromatic heterocyclyl groups comprising up to 4 hetero atoms in
20 each ring selected from oxygen, sulphur or nitrogen.

Favoured aromatic heterocyclyl groups include substituted or unsubstituted single ring aromatic heterocyclyl groups having 4 to 7 ring atoms, preferably 5 or 6 ring atoms.


In particular, the aromatic heterocyclyl group comprises 1, 2 or 3 heteroatoms, especially 1 or 2, selected from oxygen, sulphur or nitrogen.
25

Suitable values for A¹ when it represents a 5-membered aromatic heterocyclyl group include thiazolyl and oxazolyl, especially oxazolyl.


Suitable values for A¹ when it represents a 6-membered aromatic heterocyclyl group include pyridyl or pyrimidinyl.

30 Suitably R² and R³ each represent hydrogen.


Preferably, A¹ represents a moiety of formula (a), (b) or (c):



(a)




(b)



(c)

- 5 wherein: R^4 and R^5 each independently represents a hydrogen atom, an alkyl group or a substituted or unsubstituted aryl group or when R^4 and R^5 are each attached to adjacent carbon atoms, then R^4 and R^5 together with the carbon atoms to which they are attached form a benzene ring wherein each carbon atom represented by R^4 and R^5 together may be substituted or unsubstituted; and in the moiety of formula (a) X represents oxygen or sulphur.


- 10 Aptly, A^1 represents a moiety of the abovedefined formula (a).
Aptly, A^1 represents a moiety of the abovedefined formula (b).
Aptly, A^1 represents a moiety of the abovedefined formula (c).
In one favoured aspect R^4 and R^5 together represent a moiety of formula (d):



(d)

- 15 wherein R^6 and R^7 each independently represent hydrogen, halogen, substituted or unsubstituted alkyl or alkoxy.
- 20 Suitably, R^6 and R^7 each independently represent hydrogen, halogen, alkyl or alkoxy.
Favourably, R^6 represents hydrogen. Favourably, R^7 represents hydrogen.
Preferably, R^6 and R^7 both represent hydrogen.
In a further favoured aspect R^4 and R^5 each independently represent
- 25 hydrogen, alkyl or a substituted or unsubstituted phenyl group and more favourably, R^4 and R^5 each independently represent hydrogen, alkyl or phenyl.
Preferably, for the moiety of formula (a), R^4 and R^5 together represent the moiety of formula (d).
Preferably, for the moieties of formula (b) or (c), R^4 and R^5 both represent
- 30 hydrogen.
- It will be appreciated that the five substituents of A^2 include three optional substituents. Suitable optional substituents for the moiety A^2 include halogen, substituted or unsubstituted alkyl or alkoxy.

Favourably, A² represents a moiety of formula (e):




(e)

wherein R⁸ and R⁹ each independently represent hydrogen, halogen, substituted or unsubstituted alkyl or alkoxy.

Suitably, R⁸ and R⁹ each independently represent hydrogen, halogen, alkyl or alkoxy. Preferably, R⁸ and R⁹ each represent hydrogen.

Favourably, X represents oxygen. Favourably, X represents sulphur.

In one preferred aspect the present invention provides a class of compounds, which fall wholly within the scope of formula (I), of formula (II):



(II)

or a tautomeric form thereof, and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, wherein A¹, R¹, R², R³, and n are as defined in relation to formula (I) and R⁸ and R⁹ are as defined in relation to formula (e).

Suitably, n represents an integer 2, 3 or 4, notably 2 or 3 and especially 2.

Suitably, R¹ represents hydrogen, alkyl, acyl, especially acetyl, or benzyl.

When R¹ represents an alkyl group, examples of such alkyl groups include methyl and isopropyl. Preferably, R¹ represents a methyl group.

As indicated above a compound of formula (I) may exist in one of several tautomeric forms, all of which are encompassed by the present invention. It will be appreciated that the present invention encompasses all of the isomeric forms of the compounds of formula (I) and the pharmaceutically acceptable salts thereof, including any stereoisomeric forms thereof, whether as individual isomers or as mixtures of isomers.

Suitable substituents for any heterocyclyl group include up to 4 substituents selected from the group consisting of: alkyl, alkoxy, aryl and halogen or any two substituents on adjacent carbon atoms, together with the carbon atoms to which they

are attached, may form an aryl group, preferably a benzene ring, and wherein the carbon atoms of the aryl group represented by the said two substituents may themselves be substituted or unsubstituted.

When used herein the term 'aryl' includes phenyl and naphthyl optionally substituted with up to five, preferably up to three, groups selected from halogen, alkyl, phenyl, alkoxy, haloalkyl, hydroxy, amino, nitro, carboxy, alkoxycarbonyl, alkoxycarbonylalkyl, alkylcarbonyloxy, or alkylcarbonyl groups.

When used herein the term 'halogen' refers to fluorine, chlorine, bromine and iodine; preferably chlorine.

When used herein the terms 'alkyl' and 'alkoxy' relate to groups having straight or branched carbon chains, containing up to 12 carbon atoms.

When used herein the term 'acyl' includes alkylcarbonyl groups.

Suitable alkyl groups are C₁₋₁₂ alkyl groups, especially C₁₋₆ alkyl groups e.g. methyl, ethyl, n-propyl, iso-propyl, n-butyl, isobutyl or tert-butyl groups.

Suitable substituents for any alkyl group include those indicated above in relation to the term "aryl".

Suitable pharmaceutically acceptable salts include salts of the thiazolidinedione moiety, and, where appropriate, salts of carboxy groups.

Suitable pharmaceutically acceptable salts of the thiazolidinedione moiety include metal salts especially alkali metal salts such as the lithium, sodium and potassium salts.

Suitable pharmaceutically acceptable salts of carboxy groups include metal salts, such as for example aluminium, alkali metal salts such as sodium or potassium, alkaline earth metal salts such as calcium or magnesium and ammonium or substituted ammonium salts, for example those with lower alkylamines such as triethylamine, hydroxy alkylamines such as 2-hydroxyethylamine, bis-(2-hydroxyethyl)-amine or tri-(2-hydroxyethyl)-amine, cycloalkylamines such as bicyclohexylamine, or with procaine, dibenzylpiperidine, N-benzyl-b-phenethylamine, dehydroabietylamine, N,N'-bisdehydroabietylamine, glucamine, N-methylglucamine or bases of the pyridine type such as pyridine, collidine or quinoline.

Suitable pharmaceutically acceptable solvates include hydrates.

The salts and/or solvates of the compounds of formula (I) may be prepared and isolated according to conventional procedures for example sodium salts may be prepared by using sodium methoxide in methanol.

Suitable pharmaceutically acceptable salts of the thiazolidinedione moiety include metal salts especially alkali metal salts such as the lithium, sodium and potassium salts.

A preferred compound of formula (I) is 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof.

5 A compound of formula (I), or the tautomeric form thereof, and/or a pharmaceutically acceptable salt thereof, and/or a pharmaceutically acceptable solvate thereof, may be prepared using the processes described in EP 0306228. The contents of EP 0306228 are incorporated herein by reference

10 As mentioned above the compounds of the invention are indicated as having useful therapeutic properties:

A compound of formula (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, may be administered per se or, preferably, as a pharmaceutical composition also comprising a pharmaceutically acceptable carrier.

15 As used herein the term 'pharmaceutically acceptable' embraces compounds, compositions and ingredients for both human and veterinary use: for example the term 'pharmaceutically acceptable salt' embraces a veterinarily acceptable salt.

The composition may, if desired, be in the form of a pack accompanied by written or printed instructions for use.

20 Usually the pharmaceutical compositions of the present invention will be adapted for oral administration, although compositions for administration by other routes, such as by injection and percutaneous absorption are also envisaged.

Particularly suitable compositions for oral administration are unit dosage forms such as tablets and capsules. Other fixed unit dosage forms, such as powders
25 presented in sachets, may also be used.

In accordance with conventional pharmaceutical practice the carrier may comprise a diluent, filler, disintegrant, wetting agent, lubricant, colourant, flavourant or other conventional adjuvant.

Typical carriers include, for example, microcrystalline cellulose, starch,
30 sodium starch glycollate, polyvinylpyrrolidone, polyvinylpolypyrrolidone, magnesium stearate, sodium lauryl sulphate or sucrose.

Most suitably the composition will be formulated in unit dose form. Such unit dose will normally contain an amount of the active ingredient in the range of from 0.1 to 1000 mg, more usually 0.1 to 500 mg, and more especially 0.1 to 250 mg.

35 The present invention further provides a method for the treatment of atherosclerosis, in a human or non-human mammal, which comprises administering an effective, non-toxic, amount of a compound of formula (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically

acceptable solvate thereof, to a human or non-human mammal in need thereof.

The present invention also provides a method for the regulation of appetite and food intake in disorders associated with under-eating, such as anorexia nervosa, and disorders associated with over-eating, such as obesity and anorexia bulimia, in a human or non-human mammal, which comprises administering an effective, non-toxic, amount of a compound of formula (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, to a human or non-human mammal in need thereof.

Conveniently, the active ingredient may be administered as a pharmaceutical composition hereinbefore defined, and this forms a particular aspect of the present invention.


In the above mentioned treatments the compound of the general formula (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, may be taken in doses, such as those described above, one to six times a day in a manner such that the total daily dose for a 70 kg adult will generally be in the range of from 0.1 to 6000 mg, and more usually about 1 to 1500 mg.

In the treatment and/or prophylaxis of non-human mammals, especially dogs, the active ingredient may be administered by mouth, usually once or twice a day and in an amount in the range of from about 0.025 mg/kg to 25 mg/kg, for example 0.1 mg/kg to 20 mg/kg.

The therapeutic activity of a compound of formula (I), a tautomeric form thereof, a pharmaceutically acceptable salt thereof or a pharmaceutically acceptable solvate thereof may be demonstrated using conventional test methods, for example anti-atherosclerotic activity may be demonstrated using methods disclosed in Journal of Clinical Investigations 1992, Vol 89, page 706-711.

Claims

1. The use of a compound of formula (I):



(I)

- or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, wherein:


- A^1 represents a substituted or unsubstituted aromatic heterocyclyl group;
 R^1 represents a hydrogen atom, an alkyl group, an acyl group, an aralkyl group, wherein the aryl moiety may be substituted or unsubstituted, or a substituted or unsubstituted aryl group;
 R^2 and R^3 each represent hydrogen, or R^2 and R^3 together represent a bond;
 A^2 represents a benzene ring having in total up to five substituents; and
 n represents an integer in the range of from 2 to 6, for the manufacture of a medicament for use in the treatment and/or prophylaxis of atherosclerosis and/or for the regulation of appetite and food intake.

20

2. A use according to claim 1 wherein in the compound of formula (I), A^1 represents a substituted or unsubstituted, single or fused ring aromatic heterocyclyl group comprising up to 4 hetero atoms in the ring selected from oxygen, sulphur or nitrogen.

25

3. A use according to claim 1 or claim 2 wherein in the compound of formula (I), A^1 represents a moiety of formula (a), (b) or (c):



(a)

(b)

(c)


30

wherein:

R^4 and R^5 each independently represents a hydrogen atom, an alkyl group or a substituted or unsubstituted aryl group or when R^4 and R^5 are each attached to a carbon atom, then R^4 and R^5 together with the carbon atoms to which they are attached form a benzene ring wherein each carbon atom represented by R^4 and R^5 together may be substituted or unsubstituted; and in the moiety of formula (a) X represents oxygen or sulphur.

4. A use according to claim 3, wherein R^4 and R^5 each independently represent hydrogen, alkyl or a substituted or unsubstituted phenyl group.

5. A use according to claim 3 or claim 4, wherein in the compound of formula (I), R^4 and R^5 together represent a moiety of formula (d):




(d)

wherein R^6 and R^7 each independently represent hydrogen, halogen, substituted or unsubstituted alkyl or alkoxy.

6. A use according to claim 5, wherein in the moiety of formula (d) R^6 and R^7 both represent hydrogen.

7. A use according to any one of claims 1 to 6, wherein in the compound of formula (I) A^2 represents a moiety of formula (e):




(e)

wherein R^8 and R^9 each independently represent hydrogen, halogen, substituted or unsubstituted alkyl or alkoxy.

8. A use according to claim 7, wherein in the moiety of formula (e) R^8 and R^9 each represent hydrogen.

9. A use according to claim 1, wherein the compound of formula (I) is a compound
5 of formula (II):



10 or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, wherein A^1 , R^1 , R^2 , R^3 and n are as defined in relation to formula (I) in claim 1 and R^8 and R^9 are as defined in relation to formula (e) in claim 7.

15 10. A use according to any one of claims 1 to 9, wherein in the compound of formula (I) n represents an integer 2 or 3.

11. A use according to any one of claims 1 to 3, wherein in the compound of formula (I) R^1 represents a methyl group.

20

12. A use according to any one of claims 1 to 3, wherein the compound of formula (I) is selected from the list consisting of:

5-(4-[2-(N-methyl-N-(2-benzothiazolyl)amino)ethoxy]benzyl)-2,4-thiazolidinedione;

25 5-(4-[2-(N-methyl-N-(2-benzothiazolyl)amino)ethoxy]benzylidene)-2,4-thiazolidinedione;

5-(4-[2-(N-methyl-N-(2-benzoxazolyl)amino)ethoxy]benzyl)-2,4-thiazolidinedione;

5-(4-[2-(N-methyl-N-(2-benzoxazolyl)amino)ethoxy]benzylidene)-2,4-thiazolidinedione;

30 5-(4-[2-(N-methyl-N-(2-pyrimidinyl)amino)ethoxy]benzyl)-2,4-thiazolidinedione;

5-(4-[2-(N-methyl-N-(2-pyrimidinyl)amino)ethoxy]benzylidene)-2,4-thiazolidinedione;

5-(4-(2-(N-methyl-N-[2-(4,5-dimethylthiazolyl)]amino)ethoxy]benzyl)-2,4-thiazolidinedione;

- 5-(4-[2-(N-methyl-N-[2-(4,5-dimethylthiazolyl)]amino)ethoxy]benzylidene)-2,4-thiazolidinedione;
- 5-(4-[2-(N-methyl-N-(2-thiazolyl)amino)ethoxy]benzyl)-2,4-thiazolidinedione;
- 5-(4-[2-(N-methyl-N-(2-thiazolyl)amino)ethoxy]benzylidene)-2,4-thiazolidinedione;
- 5 5-[4-(2-(N-methyl-N-(2-(4-phenylthiazolyl))amino)ethoxy)benzyl]-2,4-thiazolidinedione;
- 5-(4-[2-(N-methyl-N-(2-(4-phenylthiazolyl))amino)ethoxy]benzylidene)-2,4-thiazolidinedione;
- 5-(4-[2-(N-methyl-N-[2-(4-phenyl-5-methylthiazolyl)]amino)ethoxy]benzyl)-2,4-thiazolidinedione;
- 10 5-(4-[2-(N-methyl-N-[2-(4-phenyl-5-methylthiazolyl)]amino)ethoxy]benzylidene)-2,4-thiazolidinedione;
- 5-(4-[2-(N-methyl-N-[2-(4-methyl-5-phenylthiazolyl)]amino)ethoxy]benzyl)-2,4-thiazolidinedione;
- 15 5-(4-[2-(N-methyl-N-[2-(4-methyl-5-phenylthiazolyl)]amino)ethoxy]benzylidene)-2,4-thiazolidinedione;
- 5-(4-[2-(N-methyl-N-[2-(4-methylthiazolyl)]amino)ethoxy]benzyl)-2,4-thiazolidinedione;
- 5-(4-[2-(N-methyl-N-[2-(4-methylthiazolyl)]amino)ethoxy]benzylidene)-2,4-thiazolidinedione;
- 20 5-[4-(2-(N-methyl-N-[2-(5-phenyloxazolyl)]amino)ethoxy)benzyl]-2,4-thiazolidinedione;
- 5-(4-[2-(N-methyl-N-[2-(5-phenyloxazolyl)]amino)ethoxy]benzylidene)-2,4-thiazolidinedione;
- 25 5-(4-[2-(N-methyl-N-[2-(4,5-dimethyloxazolyl)]amino)ethoxy]benzyl)-2,4-thiazolidinedione;
- 5-(4-[2-(N-methyl-N-[2-(4,5-dimethyloxazolyl)]amino)-ethoxy]benzylidene)-2,4-thiazolidinedione;
- 5-[4-(2-(2-pyrimidinylamino)ethoxy)benzyl]-2,4-thiazolidinedione;
- 30 5-[4-(2-(2-pyrimidinylamino)ethoxy)benzylidene]-2,4-thiazolidinedione;
- 5-(4-[2-(N-ccetyl-N-(2-pyrimidinyl)amino)ethoxy]benzyl)-2,4-thiazolidinedione;
- 5-(4-(2-(N-(2-benzothiazolyl)-N-benzylamino)ethoxy)benzylidene)-2,4-thiazolidinedione;
- 5-(4-(2-(N-(2-benzothiazolyl)-N-benzylamino)ethoxy)benzyl)-2,4-thiazolidinedione;
- 35 5-(4-[3-(N-methyl-N-(2-benzoxazolyl)amino)propoxy]benzyl)-2,4-thiazolidinedione;
- 5-(4-[3-(N-methyl-N-(2-benzoxazolyl)amino)propoxy]benzylidene)-2,4-thiazolidinedione;
- 5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl)-2,4-thiazolidinedione; and

5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzylidene)-2,4-thiazolidinedione; or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof.

- 5 13. A use according to claim 1, wherein the compound of formula (I) is 5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl)-2,4-thiazolidinedione; or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 94/00879A. CLASSIFICATION OF SUBJECT MATTER
IPC 5 A61K31/42 A61K31/425 A61K31/44 A61K31/505

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 5 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 419 035 (BEECHAM PLC) 27 March 1991 see page 3; claims	1-11
X	EP,A,0 306 228 (BEECHAM PLC) 8 March 1989 cited in the application see page 2; claims	1-13

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- * "A" document defining the general state of the art which is not considered to be of particular relevance
- * "E" earlier document but published on or after the international filing date
- * "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- * "O" document referring to an oral disclosure, use, exhibition or other means
- * "P" document published prior to the international filing date but later than the priority date claimed

- * "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- * "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- * "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * "&" document member of the same patent family

Date of the actual completion of the international search

7 July 1994

Date of mailing of the international search report

12.07.94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Klaver, T

INTERNATIONAL SEARCH REPORT

information on patent family members

Int: ~~Pat~~ Application No
PCT/GB 94/00879

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
EP-A-0419035	27-03-91	JP-A-	3090078	16-04-91
EP-A-0306228	08-03-89	AU-A-	2173888	09-03-89
		CA-A-	1328452	12-04-94
		JP-A-	1131169	24-05-89
		US-A-	5002953	26-03-91
		US-A-	5232925	03-08-93
		US-A-	5194443	16-03-93
		US-A-	5260445	09-11-93